

Information Summary

Organization: Neural Stem Cell Institute (NSCI) **Address:** One Discovery Drive, Rensselaer, NY 12144

Contact Person: Kathryn McIlroy **Email:** kathrynmcilroy@neuralsci.org

Phone: 518-694-8188 x 226

SCRC Code: NSCI9

Sample Name: FTD-38

Number of the Established iPSC Clones: 6

Characterization Performed:

1. Mycoplasma Test

2. Cell authentication Verification

3. Presence of the Pluripotency Marker

4. Spontaneous differentiation via embryoid body (EB) formation

5. Karyotyping

Qualified iPSC Clones: C1, C2, C3

Sample Name: NSCI9 (FTD-38)



Mycoplasma Test

Method: Determination of the mycoplasmal enzyme activity using Lonza MycoAlert Plus Detection kit.


iPSC Clone	NSCI9-C1	NSCI9-C2	NSCI9-C3
Cell passage	P6	P6	P6
B/A Ratio	0.52	0.59	0.57
Mycoplasma	NEG	NEG	NEG
Pass or Fail	Pass	Pass	Pass

Cell Authentication Verification

Markers	NSCI9 (FTD-38)	NSCI9-C1	NSCI9-C2	NSCI9-C3
D3S1358	14, 17	14, 17	14, 17	14, 17
TH01	6, 9.3	6, 9.3	6, 9.3	6, 9.3
D21S11	28, 31.2	28, 31.2	28, 31.2	28, 31.2
D18S51	12, 16	12, 16	12, 16	12, 16
Penta E	7, 15	7, 15	7, 15	7, 15
D5S818	12, 14	12, 14	12, 14	12, 14
D13S317	9, 11	9, 11	9, 11	9, 11
D7S820	8, 10	8, 10	8, 10	8, 10
D16S539	12	12	12	12
CSF1PO	10, 12	10, 12	10, 12	10, 12
Penta D	10, 13	10, 13	10, 13	10, 13
vWA	14	14	14	14
D8S1179	13, 14	13, 14	13, 14	13, 14
TPOX	8	8	8	8
FGA	23	23	23	23
AMEL	X, Y	X, Y	X, Y	X, Y
Match with original Fibroblast		Yes	Yes	Yes

Expression of the Pluripotency Markers

Antibody staining for pluripotency markers: Oct4, Nanog, Tra-1-81, SSEA-4





Differentiation Potential via EB Formation

Method: The iPSC cells were disassociated into small clumps to form EB in the low attachment dish. After a 10-day spontaneous differentiation, the EBs were harvested and RNAs were extracted. The expression levels of the selected 3-germ layer specific genes were analyzed by real-time PCR. Ct values are normalized for loading using a housekeeping gene.




Figure: Lineage specific gene expression following EB differentiation. Fold difference is shown relative to undifferentiated iPS cell.




Karyotyping

Clone: NSCI9-C1 Passage: P7

Tested: Number of metaphases counted: 20 Banding and level: GTG/550

Number of metaphases analyzed: 6 Number of karyotypes: 2

Result: 46,XY NORMAL MALE KARYOTYPE






Karyotyping

Clone: NSCI9-C2 Passage: P7

Tested: Number of metaphases counted: 20 Banding and level: GTG/550

Number of metaphases analyzed: 6 Number of karyotypes: 2

Result: 46,XY NORMAL MALE KARYOTYPE





Karyotyping

Clone: NSCI9-C3

Passage: P7

Tested: Number of metaphases counted: 20 Banding and level: GTG/550

Number of metaphases analyzed: 6 Number of karyotypes: 2

Result: 46,XY NORMAL MALE KARYOTYPE

